Attribution of Illnesses to Food Commodities

Tracy Ayers

Enteric Diseases Epidemiology Branch
Division of Foodborne, Bacterial, and Mycotic Diseases
National Center for Zoonotic, Vectorborne, and Enteric Diseases





Fundamental questions for food safety policy

>How much disease?

>Which foods?

Where does contamination occur?



Food Attribution

- Attribute the burden of foodborne diseases to specific foods
- Prioritize food safety interventions and control measures
 - What foods are causing the most human illness?



Burden of Foodborne Illness in the United States

76,000,000 cases per year Viral **Bacterial Parasitic Unknown**



Allocating foodborne disease burden: many ways to slice the pie

 Food attribution can occur at different points along the farm to fork continuum

- Can be estimated from data collected at:
 - 'Pre-harvest'
 - 'point of processing'
 - 'Point of consumption'



CDC attribution efforts

- Start with human illness data
 - > Attribute to different sources
- Approaches to Food Attribution
 - Epidemiological
 - Outbreaks and case-control studies
 - Microbiological
 - Comparison of human, animal and food subtypes



CDC epidemiological approaches to attribution

- > Outbreak data
 - > eFORS
- Case-control studies
 - > FoodNet



Using Outbreak Data for Attribution



Why use outbreak data to allocate illness by food commodity?

- For most pathogens, it is the only conclusive indication of which foods cause illness
- Represent, to varying degrees, all foodborne pathogens
- Represent a wide range of food vehicles
- Captures effect of contamination at multiple points from farm to fork



CDC Foodborne Outbreak Reporting System (eFORS)

Web-based reporting system collects epidemiological information

- Approximately 1,300 reports per year
- From 1998-2004 generated a list of nearly 2,000 implicated food items

INVESTIGATION OF A FOODBORNE OUTBREAK This form is used to report foodborne disease outbreak investigations to CDC. A boddorne authreak in defined the foodborne. The form has been participant in the control of the co				
additional info	rmation. For this investigation to be or	be counted in the CDC annual summary; Part 1 mus as much of Part 1 and Part 2 as you can.		STATE USE ONLY
	Part 1: Requ	ired Informati	ion	
Location of Exposure: Multi-state exposure Multi-state exposure Multi-county exposure Multi-county exposure Lut other states/counties in Comments, bottom of this page	Date first case became it: More Date of first known exposure: More Date of last known exposure: More More	th Day	Year Lab-confirmed Probable cases Year Estimated total	E(B)
4. Approximate Percentage Cases in Each Age Group <1 year: % 20-49 yrs: 1-4 yrs: % ≥ 50 yrs. 5-19 yrs: %	percent of total cases)% Male:%	☐ Interviews ☐ Case-cont ☐ Cohort stu	nol study plant dy Investigation a aration review (farm, marine	it factory or production it original source
7. Implicated Food(s): (based Respons listed in tarm 15 on page Could not be determined	3) hype, virulence factors, mol Etiology Confirmed* Suspected Unknown etiology Nutrale etiologies (list	Serotype t in Comments)	rasile, or toxin. If available, includence, are considered to the consideration of the consid	teristics (if avail.) that apply) (s)) cimen(s) cimen(s)
	e list on page 2, check all that app	ly)	10. Agency reporting this	outbreak:
C10	C4 C5 C6 C7 C13 C14 C15 (describe in Co- codeficial outbreaks only) P4 CP5 P6 CP7 be in Commental) NIA so only): S4 C95 (describe in Commental)	CS C9 rimenta) D N/A D P8 D P9 D N/A D No	Contact Person: NAME: TITLE: PHONE NO: FAX NO: E-MAIL: Date of completion of this Move Day Initial Report Updated Report	form:




How to use outbreak data to allocate illness by food commodity?

- Group foods in to categories that make sense for industry and regulators
- For each pathogen,
 - Sum up total number of ill
 - Calculate percent ill for each food commodity
- Finally, sum up total ill weighted by burden of illness estimates for each commodity



Hierarchical scheme for categorizing food items into commodities



Hypothetical example summing all outbreaks (not real data)

	Pei	Total U.S. foodborne				
	Beef	Pork	Veggies	Shellfis h	illnesses (CDC 1999 estimates)	
E. coli	50%	0%	40%	0%	62,458	
Vibrio	0%	0%	0%	95%	5,122	
etc						
TOTAL	%	%	%	%	14 million	



What are the challenges of using outbreak data for attribution?

- Most foodborne illness is sporadic, not associated with outbreaks
- Need sufficient number of outbreaks per pathogen
- Implicated food may not indicate implicated ingredient
 - > i.e. Complex foods



Using Sporadic Case Data for Attribution



Foodborne Disease Active Surveillance Network (FoodNet)




FoodNet Sites 2005

44.1 million (15.2% of U.S. population)



FoodNet Case-Control Studies





Driving prevention through studies of sporadic cases - Campylobacter

- ➤ FoodNet: Case-control study data intake 1998-1997
- ▶1316 confirmed cases and 1316 healthy controls (Outbreak-associated cases excluded)
- **▶**Risk Factors (Population attributable fraction):
 - Eating chicken or turkey at a restaurant (28%)
 - Eating other meat at a restaurant (21%)
 - Foreign travel (12%)
 - Contact with animal feces (6%)
 - Drinking surface water (3%)
 - Drinking raw milk (1.5%)



Blending project


e.g. E. coli O157:H7

- Summary of outbreaks*
 - 20% due to eating hamburgers
 - 4% due to animal contact
- FoodNet sporadic case-control study
 - 15% due to eating pink hamburgers
 - 8% due to visiting a farm



^{*}Rangel et al. EID 11:603-9, 2005

Blending Attribution Project





Other groups working on food attribution

- Regulatory Agencies
 - USDA, Food Safety Inspection Service
 - > FDA, CFSAN and CVM
 - Environmental Protection Agency
- Food Safety Research Consortium
 - University of Maryland
 - University of Georgia
 - University of California
 - Resources for the Future



Conclusions

- Variety of approaches used for food attribution
 - Data sources
 - Complementary approaches
- Data gaps identified
- Data sources to be used in the future
 - > PulseNet
 - > NARMS





